

Regular Vaccination Schedule

☆ Please make a vaccination schedule for your children

※ There will be no notice from the City on individual child. Please set up vaccine inoculation for your children consulting with your Doctor.

Vaccine type Inoculation method	Months (Age)																		
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
Rota virus <Rotarix Monovalent> <Rotateq pentavaent> 【Inoculation method ; Both of Rotarix and Rotateq vaccines are by oral administration.】	Over 6months(1,5mos) before 24wks(6mos)																		
HIB(Influenza b) 【Subcutaneous inoculation】	Over 2mos		①②③ of 1st series of 1st prd.						④Addition of 1st prd					before 5 yrs, (note 1.)					
Pediatric pneumococcus 【Subcutaneous inoculation】	Over 2mos		①②③ of 1st series of 1st prd.						④Add of 1st prd.					before 5 yrs, (note 1.)					
Hepatitis B 【Subcutaneous inoculation】	Over 2mos		①② of 1st series of 1st prd. ③ Add of 1st period.						Before 1yr.										
DPT-IPV (Mixture of 4kinds) 【Subcutaneous inoculation】	Over 3months			①②③ of 1st series of 1st period.						④Add of 1st					Berfore 7yrs 6mos.				
DT (Mixture of 2 kinds) 【Subcutaneous inoculation】	3mos.		1yr.						Over 1 yrs					Before 13yrs.					
BCG 【Intradermal inoculation】	5mos		group injection 4mos. Health check						before 1yr.										
MR (mixture of Measles and Rubella) 【Subcutaneous inoculation】	over 1yr.										before 2yrs.			over 5yrs,and before 7yrs, in the previous year to enter an elementary school, equivalent to senior class of a kindergarten					
Chicken pox 【Subcutaneous inoculation】	over 1 yr.										before 3yrs.								
Japanese encephalitis 【Subcutaneous inoculation】	Over 6yrs.						①,② of 1st prd.③add.					④ 2nd period			before 13yrs.				
HPV(Human Papilloma virus) 【Intramuscular injection】											Equivalent to the 6th grade of elementary school					1st grade of High school		(Note2)	

※ Contents might be changed by a revision of law, etc.

Period applicable for regular vaccination Standard regular vaccination period

Type of Vaccine : ① Live vaccine, Oral administration ② Inactivated ③ Injection of live vaccine

(Note 1: Total number of Hib vaccine and pediatric pneumococcal vaccine will depend on the timing of the 1st inoculation.

For more information, see the description from page 9.

(Note 2: HPV vaccines have different vaccination intervals between 2valent (Cervarix) and 4valent (Gardasil).

For Cervarix, 1st to 2nd is 1 months. 1st to 3rd is more than 6 months. For Gardasil, 1st and 2nd is 2 months, and 1st to 3rd is more than 6months.

[Supplements]“Inoculation period before xxx years old means that the child can be inoculated on the previous day before his or her birthday of xxx years.

Example 1 : Before one year means the child can be inoculated one the day before his or her 1st birthday.

Example 2 : In case of a child born on Feb.29, the previous day of his or her birthday is Feb. 27 in the years without Feb. 29.